

A Comprehensive Review in Novel Therapeutics in Acute Leukemia

Amir Gholamzad¹, Ghazaleh Daryoushi¹, Mahsa Khatibi¹, Sajad Khonche², Arvin Ahmadpour², Matin Malek², Arya Mehrkhah², Haniyeh Rezazadeh¹, Arghavan Bazrafshan Sichani², Parasto Jafari², Kimiya Shahabi², Fatemeh Ezzati², Seyed Ali Hosseini Zavareh^{3,*}

¹Medical Laboratory Sciences Student, Faculty of Allied Medicine, Tehran Medical Sciences, Islamic Azad University, Tehran, Iran

²Medical Student, Faculty of Medicine, Tehran Medical Sciences, Islamic Azad University, Iran

³Medical Student, Department of Medicine, Islamic Azad University Tehran Medical Sciences, Iran

*Corresponding author: Seyed Ali Hosseini Zavareh, Department of Medicine, Islamic Azad University Tehran Medical Sciences, Iran

Received: April 21, 2022

Published: May 16, 2022

Abstract

In recent years, acute leukemia has become a challenging disease for children, adolescents and adults, so there are always different treatments; None of them are considered as a definitive treatment for this disease. Finding treatments that have fewer side effects and more results has always been one of the most important and recently discussed challenges. The production of vaccines for the prevention and treatment of cancer has always been one of the candidates for the treatment of various cancers, and recently the companies Estraznika and Pfizer have achieved very good results. These days, new therapies called gene therapy have been introduced, which have shown promising results. T cell therapy is one of these treatments that has shown very good results and is currently undergoing clinical phases in various cancers, especially acute leukemias, in the United States of America. On the other hand, science-based therapy Virology is underway as virus therapy, which is undergoing animal testing and clinical trials for various cancers and viruses. It is hoped that viruses can be used as allies in the fight against cancer. With the help of microbiology, this problem has reached acceptable results and they have been able to study the microbiomes of leukemia patients and find out the results that contribute to a successful chemotherapy treatment. In this article, we aim to explore new treatments for acute leukemia and examine the results of different trials and different clinical phases. It is hoped that with the advancement of science, the treatment of leukemia patients will always be possible.

Keywords: Acute Leukemia; Cancer vaccine; Virotherapy; Nanomedicine therapy; Immunotherapy

Introduction

Acute Leukemia (AL) is a type of leukemia that originates from the Bone Marrow (BM). Due to the abnormally rapid growth of white blood cells, patients often experience fatigue, bleeding, bruising, fever, and a weakened immune system. According to lineage, AL can be classified as Acute Lymphocytic Leukemia (B) (B-ALL), acute T-Cell Lymphocytic Leukemia (T-ALL), and Acute Myelogenous Leukemia (AML). Leads to bone marrow disorders Most patients experience fatigue, bleeding, bruising, and weakened immune systems due to the abnormally rapid growth of white blood cells Acute leukemia can be classified as acute B-cell lymphocytic leukemia, leukemia Acute T-cell lymphocytes and ultimately acute myelogenous leukemia Acute lymphocytic leukemia is more common in children, while acute myeloid leukemia is a more common type of acute leukemia in the elderly [1-5] CAR T Cell Therapy is one of the Recent treatments for acute leukemia have been in the clinical stages for other cancers and have produced significant clinical responses to specific subsets of leukemia or B-cell lymphoma, but there are limitations. And barriers such as tumor resistance to CAR structures are still problematic [6].

This acute leukemia has also been used. Nano-based targeted therapies have also been used for various diseases, including cancer, and various drugs have completed or completed their clinical phase, but these therapies are known as complementary therapies; Therefore, considering the mechanisms of suppression of the immune system as a contributing factor in treatment, combining cancer vaccines with other immune therapies, or in other words, the use of complementary therapies for the treatment of cancer, seems to be more effective [7]. Has been proposed for target therapy; Oncolytic virus therapy is one of these therapies that tests its trials and clinical phases for various viruses and cancers and faces challenges such as not getting infected with the virus because according to the principles of oncolytic virus as a virus Genetically engineered is defined and, unlike the gene therapy itself, the virus is used, which can selectively multiply in cancer cells and kill them without damaging normal tissues [8].

But apart from the new therapies, the current therapies also have challenges. One of these challenges is to reduce the side effects of the treatments, which has been made possible by examining the intestinal microbiome.

Nano AML

Therapeutic management of AML disease involves the induction of chemotherapy and the recovery phase. Higher invasive treatment, such as allogeneic bone marrow transplantation, is performed for high-risk patients. Recommended treatments for patients in induction failure, clinical trials, low-intensity treatment, azacitidine (HCT, allogeneic decitabine, or supportive therapy) [9] Most AML patients are over 60 years old, and many receive light supportive or chemotherapy. The reason for the toxicity and unclear efficacy of invasive treatment is recommended. The 5-day low-dose treatment with Decitabine has the same outcome as the initial treatment in elderly AML patients, and its acceptable toxicity and mortality are 30 days [10,11]. The combination of talacotuzumab and decitabine is less beneficial than decitabine alone in elderly patients with AML [12]. The monosomal and complex karyotypes were Azacitidine, which saw a 46-46% reduction in the risk of death in patients treated with this drug compared to patients with conventional care regimens [13,14], the rate of MRD in 80% of patients after receiving 4 courses Azacitidine therapy, in the absence of relapse Hematologic, decreased or remained constant. Even with azacitidine re-treatment in most patients, hematologic relapses were unavoidable [101006]. Treatment with azacitidine for 32 days for 32 days is safe and can also be used for at least 4 periods after allogeneic transplantation for AML / MDS patients undergoing severe treatment [9]. We can also see more benefits after more cycles [17]. In addition, nanotechnology has made the treatment of many cancers, such as acute myeloblastic leukemia (AML), easier and more accurate than the above methods. Three nanotechnology methods can be used to treat blood malignancies, which are: 1. Organic nanoparticles such as dendrimers, polymer particles and Lipid Based NPs. Inorganic nanoparticles such as iron particles, QDs, carbon-based particles and mesoporous silica. Nano-hybrid [18,19], for the treatment of Acute Myeloblastic Leukemia (AML) there are two chemotherapy and immunological methods [20]. Organic nano method can be used for both diagnosis and treatment of this disease, which is combined with the above two methods. The only disadvantage of using Nanomedicine is that it causes small subcellular damage to organ damage [21]. In this cancer, abnormal blood cells multiply uncontrollably and can invade and infect other organs. Using nanocarriers, anti-cancer drugs can be targeted into infected tissues and cells, and by increasing the half-life of the drug, reducing the dosage form, and minimizing drug interactions and drug toxicity, the therapeutic efficacy can be specifically addressed. Maximized [22,23]. In one study, a comparison was made between treating AML patients who did not qualify for invasive chemotherapy, between the effect of a single drug and the same drug with a nanoparticle. The incidence of fibril neutropenia was also much higher than in patients treated with Azacitin alone [24]; As a result, AML management methods with ventoclus-based compounds are expected to evolve and be considered more seriously over the next few years [25]. For this reason, the production of a precursor from nanoparticles called -CD-44targeted glutathione-sensitive hyaluronic acid mercaptopourin was provided. According to the evidence obtained, this prodrug of nanoparticles has very low side effects as well as high suppressive power of OCI / AML2 tumors in leukemia, which has made it a very functional and effective drug nanoparticle [26]. In another study, it was observed that the two molecules CD44 and NF-KB were much more expressed in leukemia patients than

in normal people (25 times) [18].

For the treatment of leukemia patients, an encapsulated PTL nanoparticle called nano-antiCD44 encapsulating PTL was designed which acted on the NF-KB molecule and reduced cell proliferation in this molecule by 50% and was very effective in treating the disease [27]. Another drug that is very effective in treating AML, called Vorinostat, was tested and found to inhibit the activity of HDAC class 1 and 2 enzymes at nanomolar concentrations and induce cytotoxic activity at micromolar concentrations. A protein-based nanomedicine from Vorinostat encapsulated by human serum albumin, which increases the solubility and enhances the effect of the drug, and the combination of this drug with DNMTi increases the therapeutic efficacy [28,29]

CAR T cell Therapy AML

AML is a heterogeneous blood malignancy that often has a poor prognosis and has the highest annual death rate from leukemia. High levels of CD123 and CD33 have been observed in these patients, with CD123 being the most common antigen in AML [30,31,32]. But despite numerous treatments for AML, there are still refractory AMLs. [33] In AML cells, the 13-CD protein is highly expressed, targeting cancer cells Strongly destroyed. But direct targeting of CD13 can cause toxicity to human HSCS and other normal cells. [34] However, the use of CAR T + CD19 (which is effective in treating CD19 + B cell leukemia) can limit the expression of CD13, CD19, and CD123 proteins in cancer cells without having cytotoxicity. Studies have shown that concomitant use of SB vectors with CAR T not only increases the effectiveness of CAR T, but also reduces costs. However, there are limitations for SB, including low gene transfer and toxicity to T cells. [35,36,37] In addition, the use of ScFv also affects CAR T cytotoxicity [38]. Most CAR T cells today use anti-human SCFV extracted from mice [39]. One of the problems in treating AML with CAR T is the lack of appropriate antigen. Because Flt3Lg stimulates cell proliferation in AML, disruption of the Flt3Lg binding site in Flt3 may inhibit blast cell proliferation [40]. More generally, problems with AML treatment with CAR T include cases such as Antigenic Heterogeneity and On-Target Off-Tumor Toxicity [41]. GMP was examined there and showed higher sensitivity to FLT3 CAR T cells in the laboratory. Therefore, to prevent cytotoxicity, FLT3 CAR can be modified by having an induction-switching switch with retoximab. Or after AML eradication, reduce FLT3 CAR cells to recover bone marrow [42]. Molecule 1 - CLL is present on the surface of AML cells but not on the surface of normal HSCS cells, so it is considered as a safe antigen for CAR T to kill AML cells as well as LSCS. [36] No toxicity has been reported for CAR T against 1 CLL for CD34 and CD38 + cells. CLL is expressed in myeloid cells, so CAR anti-1 CLL cells effectively destroyed AML blasts so that they had the same toxicity as HL60 under the same conditions. Overall, CAR T against 1 CLL shows excellent efficacy in the treatment of AML in vivo and has no life-threatening side effects [43]. Research has also shown that CD28 / CD27 excitatory signals can reduce the persistence of CAR T against CLL1 [44]. UniCAR T in combination with TM123 has been shown to be effective in killing leukemia + CD123 cells and has been shown to reduce side effects. Treatment of HSPCs with UniCAR-T guided by TM123 resulted in lysis of HSPCs expressing CD123 in the laboratory, but this is transient and canceled by removal of TM123. The combina-

tion of CAR T anti-CD123 with AZA has also been shown to be effective in the treatment of AML without tissue damage and hematopoietic insufficiency [45,46]. CD7-specific CAR T cells effectively invade CD7-expressing AML cells without cytotoxicity to normal myeloid and erythroid cells. In fact, CD7 CAR Ts are a transient treatment and provide the conditions for stem cell transplantation, which terminates CAR T activity and resets NK cells and T lymphocytes [47]. Finally, it can be noted that despite advances in the use of T CAR cell therapy in this disease, AML has been obscure to some extent due to the lack of truly specific surface antigens [48].

Virotherapy AML

In blood diseases treated with systemic OVT, virus-mediated immunotherapy is performed, which includes tumor cells and responsive immune cells. The first injection of CVA21 (which shows anti-tumor responses in blood diseases) stimulated ISG and increased the expression of CD69 (CD69 on lymph nodes, causing cell migration and uptake of nutrients and preservation of lymphocytes) on CD4 + T cells. NK and TCD8 + became on day 3 but decreased until day 22. CVA21 showed a high capacity for immunotherapy responses that, through ICAM-1 and dendritic plasmacytoid cells, was able to activate NKs, which could be a lethal pathway for malignant cells. Combining CVA21 with other immunotherapies and injecting it seems to be effective in treating AML and MM [49]. Another case in point is Reovirus, an RNA virus that mainly signals cells Hyperactivity, for example, infects the Ras pathway. In using the reovirus for extracorporeal clearance, MM cells were incompletely cleaved and only 50% of the MM cells were effectively cleaved. There were no follicular and burkit lymphoma cells. Therefore, in conclusion, reovirus may have therapeutic potential in some hematologic malignancies, but its efficacy should be evaluated in further clinical trials [50].

Vaccine AML

Immune cells are used to identify tumor cells that are difficult to target because of the immune mechanism. Based on the efforts made and considering the complexity of immune suppression mechanisms, the use of combination therapy as a combination of cancer vaccines with other immunotherapy seems to be more effective [51] in a vaccine study aimed at cells NKT and their combination with anti-4-1BB resulted in the survival of all mice with MLL AML because the vaccine in combination with anti-4-1BB activates CD8 + T and produces IFN- γ , which clears the tumor. It is done by increasing CD8 + T and IFN γ . But after combination treatment of 1-anti-PD and anti-4-1BB, 40-60% of mice survived. Vaccine and anti-4-1BB alone, either in combination or 1-anti-PD, increased AML tumor clearance in some mice, but the tumor gradually grew in some mice that did not respond to the vaccine. Evidence has shown that delays in the use of 1-anti-PD can counteract the therapeutic effect. Therapeutic response to NKT and anti-4-1BB vaccination cleared the tumor in all affected organs of infected mice, including blood, bone marrow and spleen [52,53]. A clinical trial using SVF cells incubated with vaccine virus It was performed to treat 26 patients with AML, which showed that the combination of SVF and ACAM2000 was safe, and that inflammatory symptoms associated with the virus were seen in most patients within 2 weeks of treatment. Skin rash and viral DNA survival in the blood were among the signs of vaccine virus survival in the body after treatment. In some patients, the combination of this virus with inhibition

of checkpoints caused the death of tumors, but it should still be evaluated in a larger population [54,55]. In one study, a measles vaccine was used to treat AML patients. Leukemic blasts in patients were lysed by MeV, and although they were susceptible to the virus, CD46 levels were either normal or decreased. Addition of 5-FC enhances the tumorigenic effects of MeV-SCD but is less commonly used in treatment due to its gastrointestinal toxicity and low potency. [56]. IL-15 DCS has been shown to be able to inhibit 75 T cells. Which indicates its anti-tumor activity with 15-IL secretion. The DC vaccine in combination with anti-1-PD drugs further affects AML cells. T 75 cells in these patients are susceptible to stimulation of 15-IL DC, especially in the presence of IPP. Unlike aB T cells, T 75 cells are not dependent on the tissue compatibility complex and antigen-presenting cells [57,58]. The combination of IL-15 / IL-15Ra arteriodimer with CD80 in autologous leukemic cells as a vaccine seems to be an effective treatment for AML and expression of 15-IL in AML vaccines stimulates immunity and reduces toxicity. It becomes a cell. Mice with more severe AML responded to the 32Dp210-IL-15 / IL-15Ra / CD80 vaccine, which allowed them to survive longer and eradicate leukemia [59,60]. The DC vaccine targets WT1 with the ability to increase the immunity of non-leukemic T cells in AML patients, and CD8 + cells with WT1 remove the remaining tumor cells and reduce the recurrence of AML [61]. WT1 vaccination was performed on 22 AML patients which was effective and no mortality was reported [62,63].

In a phase safety test DCP-001 vaccination was evaluated in patients with advanced AML. MRD after chemotherapy is a major challenge in the treatment of AML. This vaccine has the ability to stimulate cellular and humoral immunity. The vaccine was injected every 4 weeks and 2 weeks, and the result was that elderly patients with stable and adjusted immune systems showed a greater response to the vaccine. "DCP-001 vaccination was effective in patients with safe elderly AML. In patients with CR and stable levels of blood T cells, it caused long-term survival (36 months). Therefore, it can be an effective therapeutic approach in the treatment of AML for complete recovery after chemotherapy [64]. Therefore, in an article, the use of hUC - MSCS as a carrier for reoviruses in the treatment of AML was investigated. Reovirus infection caused the secretion of inflammatory cytokines from hUC - MSCS, especially CXCL10, which increased the antitumor effect of the virus. In fact, hUC-MSCS prevented the reovirus from being neutralized by high-titer antiviral antibodies and also facilitated the transmission of the reovirus to tumor cells. But in general, the role of hUC - MSCS as an antitumor needs further study. MSCS, along with reovirus, infiltrated lymphocytes into tumor cells and reduced immunosuppression. According to the results, reovirus activated AKT and SAPK / JNK and also increased the expression of CXCR4 and CXCR7 in hUC - MSCS and it seems that AKT plays an important role in cell migration and virus replication [65,66] in an experiment with a nanoparticle vaccine coated with an antigen-rich AML cell membrane (AMCNP) has been investigated to treat tumors and improve inhibition of cancer recurrence. The vaccine also protects against AML and cancer-specific T cell responses. Increase blood. It has more advantages than WCL vaccines. In the bone marrow and liver of mice vaccinated with AMCNP, much fewer leukemia cells were observed than in mice vaccinated with WCL. The AMCNP vaccine made here is multi-antigen and fully personalized, eliminating the need to identify neo-antigens. These vaccines can be combined with a variety

of adjuvants [67,68].

MRD 9 fact, TRAIL-coated ZA4-ovcovirus adenovirus -SZU vaccine (to enhance tumor targeting ability) showed strong antitumor effect against AML as well as combination zA4 with Rh2 treatment of leukemia and BALB / C cell transplantation Increased mice [30]. In one experiment, tumor cells were treated with DACA and a new conjugate TLR7 agonist 106-SZU. Cells conjugated to 106-SZU relative to DAC perform DC maturation and T activity in vitro against tumor cells and inhibit them. That is, combining the DAC with 106-SZU, in addition to activating TLR7, has the ability to deliver tumor antigen to DCs to remove tumors by binding TLR7 to the DC surface. DAC-AML-106 shows potential therapeutic effect in combination with PDI or PDL1 monoclonal antibody [69]. Combining VSV virus with CAR T showed that T cells have the ability to load VSV and protect it against the immune system have. Experiments on NSG mice showed that loading high doses of VSV virus on CD8 + T cm had a stronger anti-leukemic effect than VSV alone. Therefore, the combination of an oncolytic virus with effective human tumor-specific immune cells as virus carriers could have a significant therapeutic effect compared to monotherapy [70]. It can be noted that the development of nanotechnologies in recent years has provided potential strategies to improve cancer immunotherapy, and its important advantages, as mentioned earlier, include the precise targeting of specific cell types in the design of cancer vaccines and Delivery of safety modifiers used [71].

CAR t - cell therapy for ALL

The treatment of patients with Acute Lymphoblastic Leukemia (ALL) using CAR - T cells as a new approach has been considered by many researchers. Among the numerous CAR - T receptors, CD19 receptor has been studied as an effective receptor for CAR - T, especially in the treatment of patients with recurrent B - ALL patients after transplantation with poor prognosis [72,73]. Anti-CD19 CAR-T is rare and non-hematologic diseases such as MDS, hypogammaglobulinemia and hypoglycemia have been observed in some cases [74]. In a report, mild neurotoxicity with lower degrees and severe neurotoxicity with higher degrees was observed for -Anti CD19 CAR-T, with factors such as pre-treatment disease level and stimulatory agonists such as N-methyl-D-aspartate (NMDA) receptor quinoa. Acid and glutamate in bone marrow are involved in neurotoxicity [75]. In patients with Relapsed / Refractory Acute (r / rALL Lymphoblastic Leukemia) and Non-Hodgkin Lymphoma (NHL) treated with CAR T, cases of CMV and RSV infection with pneumonia were reported, which did not pose a serious risk because most of these infections. It was observed in the first days of CAR T injection and in hospital conditions [76]. CRS (Cytokine release syndrome) has also been reported as a result of effective treatment of recurrent B - ALL individuals after transplantation with CAR T cells [77]. The use of blinatumomab before CAR T treatment reduces the recovery and development of MRD, thereby reducing the effect of CAR T on B - ALL treatment [78,79] AUTO3 as a type of CAR T designed to target CD19 and CD22 has been shown to be effective in the treatment of children with r / r BALL simultaneously with a dose of 106 CAR T - cells 23 Kg. Although the use of AUTO3 has side effects such as neutropenia, anemia, thrombocytopenia, fever, but causes Complete Recovery (CR) or minimal residual disease (MRD) [80]. Also, treatment with CD22 CAR T after treatment with CD19 CAR T increases the function of CD19 CAR T and the half-life of CAR T, so the use

of CD19 and CD22 can show potential effects in the treatment of patients B - ALL [81] KTE - X19 to As a type of autologous CAR T CD1919, in the treatment of r / r B - ALL patients despite complications such as fever (42%), hypotension (40%), decreased platelet count (33%), anemia (31%) Hypophosphatemia (31%), hypoxia (24%), encephalopathy (22%), neutropenia fever (22%) and a decrease in neutrophil count (22%) in some patients result in complete or minimal remission of the disease [82 The other two receptors for CAR T, including CD28 and 4-1BB, show an antitumor response in ALL patients with mild side effects, but these two have different response patterns [83].

Vaccine for ALL

Potential for ALL treatment. One of the vaccines that has the ability to activate the immune system and kill tumor cells without side effects is the IPSC-based vaccine. This vaccine demonstrates anti-tumor effects against T-ALL by delivering it to T cells via DCs. [84]. For example, the use of Epstein-Barr virus-specific cytotoxic T cells (EBV CTL) can reduce neurotoxicity and CRS (cytokine release syndrome) and GVHD due to CAR T injection, thereby increasing the persistence of CAR Ts in the body [85]. CAR T cells with the epidermal growth factor receptor pathway substrate vaccine 8 (Eps8 - DCS peptide - derived dendritic cell), reduce cell death due to CAR T injection and its side effects, thus increasing its shelf life in the body. This vaccine induces anti-tumor effects by increasing the secretion of 2-IL and TNF- α and increases the proliferation of T cells and the persistence of CAR T in the body, thus overcoming recurrent leukemia cells [86].

Nano medicine for ALL

A nanoparticle called novel CHC-SDS was made, which increased the hydrophobicity of the SDS fraction and also increased the pyrazoline H3Tm04 without encapsulation [87,88]. According to experiments, these nanoparticles are compatible with the living and non-toxic tissue environment, and CHC-SDS nanocapsules are a very good delivery system for a variety of pyrazoline-derived and hydrophobic drugs [87]. In one experiment, 737-ABT and IRAK1 / 4 inhibitors were inserted into a capsule of Polyethylene glycol modified poly nanoparticles, which had a double effect in inducing T-ALL apoptosis compared to the combination of IRAK1 / 4 and AB37-737 solution. Also, IC50 for -IRAK / ABT NPS was twice as low as for free combination drugs in Jurkat cells [89]. In another experiment, the effect of active thiol polymers on cell relationship and nanoparticle adsorption was studied and it was found that thiol-reactive star polymers contain mPEG brush corona, which contains some diethylene glycol brush moieties that improve cells. Leukemia becomes delirium [90]. Another nanoparticle that is very suitable for the treatment of ALL was called BSA / ASN25% / Pol407, which performed very well in intravenous injection and protected ammonium against free enzymes, which prevents hyperammonemia [91,92]. A new therapeutic strategy against ALL was the fabrication of DOX-PMs-NPMBP nanoparticles, which significantly inhibited tumor growth in ALL and also showed no systematic toxicity after treatment [93]. One of the markers in the malignant membrane of malignant cells, including B - LL, is CD19, and a nanoparticle called CD19 - PEG - MTN / DOX was made, which forced the CD19 + B - LL cell into the apoptor, but Not toxic to normal cells. After treatment with DOX / nanoparticle, CD19 - PEG - MTN Bax 6 - and NALM post-apoptotic pro-

teins were dramatically upregulated and anti-apoptotic proteins such as 70 Bcl2, MCL - 1, HSP, were downregulated. Indicating the activation of the apoptotic pathway by nanomedicines [94].

Virotherapy for ALL

Virus therapy for leukemia is still in need of further research; But this treatment, along with other treatments, has had good results. In a simultaneous study using rituximab, oncolytic measles virus was used, which boosted the immune system; On the other hand, increasing the effectiveness of oncolytic viruses has become one of the challenges that currently the use of bone marrow stem cells as carriers of oncolytic virus and the use of antibodies against it to prevent toxicity is one of the best methods. Measles was contracted here from the oncolytic virus. [95,96].

Microbiome AML, ALL

Intestinal barrier is a vulnerable site in patients who have received intensive chemotherapy and in patients with hematologic malignancies, intestinal microbiota disorder has been well established with chemotherapy and broad-spectrum antibiotics [97]. Maintaining high microbial diversity and metabolites produced by these microbiotas in the intestine helps reduce toxicity and initiation of chemotherapy reduces this diversity and several complications [98,99]. For example, short-chain fatty acids (SCFAS) are microbial metabolites that are derived from dietary fiber in the colon and play an important role in body health. Butyrate is one of the most important SCFASs, which is the epithelial food source and reduces the inflammatory response of the intestine by inhibiting the production of cytokines, and Lachnospiraceae species, which are usually part of the intestinal microbiota and produce butyrate, are reduced by starting chemotherapy [100,101]. Indole compounds, which have antioxidant and anti-inflammatory effects, are also produced from tryptophan metabolites by intestinal microbiota, especially Commensal clostridium and *E. coli*, which are significantly reduced in patients with acute leukemia undergoing induced chemotherapy [102]. Overgrowth of Enterococcaeae species is also seen at the start of chemotherapy, which may cause acute GVHD and ulcerative colitis by compromising intestinal epithelial integrity and stimulating macrophage activation. In the past, an increase in the incidence of enterococci has been reported as a strong predictor of infectious complications in ALL children and adult AML [98,99]. In addition, the spread of Akkermansia species in the gut increases the risk of neutropenia (NF) in patients with acute leukemia receiving intensive chemotherapy, and adult AML patients with the first febrile neutropenia after intensive chemotherapy have a significant reduction in microbiota diversity. Intestines show [99,102]. Bloodstream Infection (BSI) pathogens are said to be the leading cause of death in ALL patients, and microbiota is a major source of BSI-causing pathogens in immunocompromised patients, and functional changes in intestinal bacteria in ALL patients may be possible. Increase BSI [101].

AFMT (Autologous Stool Microbiota Transfer) is one of the methods of intestinal microbial repair in patients with acute leukemia, after AFMT in AML patients who underwent chemotherapy and broad-spectrum antibiotics. Ruminococcaeae were observed. However, further research has shown that allogeneic FMT (fecal matter transferred from a healthy person to a person with intestinal microbiome disorder to repair and regenerate diseased microbiome) is safer than AFMT, and that the use of FMT is safer. It is effective in hematological malignancies for reducing infections related to chemotherapy and

hematopoietic stem cell transplantation [103,104]. In general, intestinal microbiota play an important role in the recovery status of patients with acute leukemia [105].

References

1. Dorez GM, Devesa SS, Curtis RE, Linet MS, Morton LM. "Acute leukemia incidence and patient survival among children and adults in the United States." 2001–2007. *Blood*, 2012; 119: 34–43.
2. Terwilliger T, Abdul-Hay M. "Acute lymphoblastic leukemia: A comprehensive review and 2017 update." *Blood Cancer J*, 2017; 7: e577.
3. Dohner H, Estey E, Grimwade D, Amadori S, Appelbaum FR, Buchner T, et al. "Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel." *Blood*, 2017; 129: 424–447.
4. Guru Murthy GS, Pondaiah SK, Abedin S, Atallah E. "Incidence and survival of T-cell acute lymphoblastic leukemia in the United State." *Leuk. Lymphoma*, 2019; 60: 1171–1178.
5. Dohner H, Weisdorf DJ, Bloomfield CD. "Acute Myeloid Leukemia." *N. Engl. J. Med*, 2015; 373: 1136–1152.
6. Sterner Robert C, Rosalie M Sterner. "CAR-T cell therapy: Current limitations and potential strategies." *Blood cancer journal*, 2021; 11(4): 1-11.
7. Wu Ming, et al. "Therapeutic cancer vaccine therapy for acute myeloid leukemia." *Immunotherapy*, 2021; 13(10): 863-877.
8. Fukuhara, Hiroshi, Yasushi Ino, Tomoki Todo. "Oncolytic virus therapy: A new era of cancer treatment at dawn." *Cancer science*, 2016; 107(10): 1373-1379.
9. Tallman Martin S, et al. "Acute myeloid leukemia, version 3.2019, NCCN clinical practice guidelines in oncology." *Journal of the National Comprehensive Cancer Network*, 2019; 17(6): 721-749.
10. Cashen Amanda F, et al. "Multicenter, phase II study of decitabine for the first-line treatment of older patients with acute myeloid leukemia." *Journal of Clinical Oncology*, 2010; 28(4): 556-561.
11. Hiddemann Wolfgang, et al. "Management of acute myeloid leukemia in elderly patients." *Journal of Clinical Oncology*, 1999; 17(11): 3569-3576.
12. Montesinos Pau, et al. "Safety and efficacy of talacotuzumab plus decitabine or decitabine alone in patients with acute myeloid leukemia not eligible for chemotherapy: results from a multicenter, randomized, phase 2/3 study." *Leukemia*, 2021; 35(1): 62-74.
13. Döhner Hartmut, et al. "Cytogenetics and gene mutations influence survival in older patients with acute myeloid leukemia treated with azacitidine or conventional care." *Leukemia*, 2018; 32(12): 2546-2557.
14. Wen Bingbing, et al. "Indirect comparison of azacitidine and decitabine for the therapy of elderly patients with acute myeloid leukemia: A systematic review and network meta-analysis." *Experimental hematology & oncology*, 2020; 9(1): 1-10.
15. Platzbecker U, et al. "Azacitidine for treatment of imminent relapse in MDS or AML patients after allogeneic HSCT: results of the RELAZA trial." *Leukemia*, 2012; 26(3): 381-389.
16. Schuh Andre C, et al. "Azacitidine in adult patients with acute myeloid leukemia." *Critical reviews in oncology/hematology*, 2017; 116: 159-177.
17. De Lima Marcos, et al. "Maintenance therapy with low-dose azacitidine after allogeneic hematopoietic stem cell transplantation for recurrent acute myelogenous leukemia or myelodysplastic syndrome: a dose and schedule finding study." *Cancer*, 2010; 116(23): 5420-5431.
18. Huang Lifan, et al. "Nanomedicine—a promising therapy for hematological malignancies." *Biomaterial's science*, 2020; 8(9): 2376-2393.
19. Ansari Mohammed T, et al. "Lipid-based nanocarriers for cancer and tumor treatment." *Current pharmaceutical design*, 2020; 26(34): 4272-4276.
20. Grosso Dolores A, Rosemary C Hess, Mark A Weiss. "Immunotherapy in acute myeloid leukemia." *Cancer*, 2015; 121(16): 2689-2704.
21. Tatar Andra-Sorina, et al. "Nanomedicine approaches in acute lymphoblastic leukemia." *Journal of Controlled Release*, 2016;

- 238: 123-138.
22. Sauvage Félix, et al. "The use of nanocarriers in acute myeloid leukaemia therapy: challenges and current status." *Current Pharmaceutical Biotechnology*, 2016; 17(1): 30-41.
 23. DiNardo, Courtney D, et al. "Azacitidine and venetoclax in previously untreated acute myeloid leukemia." *New England Journal of Medicine*, 2020; 383(7): 617-629.
 24. Othman Tamer A, et al. "An evaluation of venetoclax in combination with azacitidine, decitabine, or low-dose cytarabine as therapy for acute myeloid leukemia." *Expert review of hematology*, 2021; 14(5): 407-417.
 25. Qiu Jie, et al. "Glutathione-sensitive hyaluronic acid-mercaptopurine prodrug linked via carbonyl vinyl sulfide: a robust and CD44-targeted nanomedicine for leukemia." *Biomacromolecules*, 2017; 18(10): 3207-3214.
 26. Sami Shaheedul A, et al. "Current and future molecular targets for acute myeloid leukemia therapy." *Current Treatment Options in Oncology*, 2020; 21(1): 1-16.
 27. Chandran Parwathy, et al. "Epigenetics targeted protein-vorinostat nanomedicine inducing apoptosis in heterogeneous population of primary acute myeloid leukemia cells including refractory and relapsed cases." *Nanomedicine: Nanotechnology, Biology and Medicine*, 2014; 10(4): 721-732.
 28. Prebet Thomas, Norbert Vey. "Vorinostat in acute myeloid leukemia and myelodysplastic syndromes." *Expert opinion on investigational drugs* 2011; 20(2): 287-295.
 29. Baroni ML, Martinez DS, Aguera FG, Ho HR, Castella M, Zanetti SR, et al. "41BB-based and CD28-based CD123-redirection T-cells ablate human normal hematopoiesis in vivo." *Journal for immunotherapy of cancer*, 2020; 8(1).
 30. Liu J, Tong J, Yang H. "Targeting CD33 for acute myeloid leukemia therapy." *BMC cancer*, 2022; 22(1): 1-7.
 31. Tatsumi Eiji. "A mini-review of CD13 antigen in AML: easy induction or enhancement of expression in in vitro culture and necessary consideration for assessment." *The Southeast Asian Journal of Tropical Medicine and Public Health*, 2002; 33: 155-157.
 32. Meyer JE, Loff S, Dietrich J, Spehr J, Jurado Jiménez G, von Bonin M, et al. "Evaluation of switch-mediated costimulation in trans on universal CAR-T cells (UniCAR) targeting CD123-positive AML." *Oncoimmunology*, 2021; 10(1): 1945804.
 33. He X, Feng Z, Ma J, Ling S, Cao Y, Gurung B, et al. "Bispecific and split CAR T cells targeting CD13 and TIM3 eradicate acute myeloid leukemia." *Blood*, 2020; 135(10): 713-723.
 34. Rotiroti MC, Buracchi C, Arcangeli S, Galimberti S, Valsecchi MG, Perriello VM, et al. "Targeting CD33 in chemoresistant AML patient-derived xenografts by CAR-CIK cells modified with an improved SB transposon system." *Molecular Therapy*. 2020; 28(9): 1974-1986.
 35. Wang J, Chen S, Xiao W, Li W, Wang L, Yang S, et al. "CAR-T cells targeting CLL-1 as an approach to treat acute myeloid leukemia." *Journal of hematology & oncology*, 2018; 11(1): 1-3.
 36. Hofmann Susanne, et al. "Chimeric antigen receptor (CAR) T cell therapy in acute myeloid leukemia (AML)." *Journal of clinical medicine*, 2019; 8(2): 200.
 37. Chitre S, Gaken J, Venuso A, Mufti GJ. "Single and dual targeting chimeric antigen receptor T-cell therapy in acute myeloid leukemia." *Blood*, 2020; 136: 25.
 38. John S, Chen H, Deng M, Gui X, Wu G, Chen W, et al. "A novel anti-LILRB4 CAR-T cell for the treatment of monocytic AML." *Molecular Therapy*, 2018; 26(10): 2487-2495.
 39. Maiorova V, Mollaev MD, Vikhрева P, Kulakovskaya E, Pershin D, Chudakov DM, et al. "Natural Flt3Lg-Based Chimeric Antigen Receptor (Flt3-CAR) T Cells Successfully Target Flt3 on AML Cell Lines." *Vaccines*, 2021; 9(11): 1238.
 40. Fiorenza Salvatore, Cameron J Turtle. "CAR-T cell therapy for acute myeloid leukemia: preclinical rationale, current clinical progress, and barriers to success." *BioDrugs*, 2021; 35(3): 281-302.
 41. Sommer C, Cheng HY, Nguyen D, Dettling D, Yeung YA, Sutton J, et al. "Allogeneic FLT3 CAR T cells with an off-switch exhibit potent activity against AML and can be depleted to expedite bone marrow recovery." *Molecular Therapy*, 2020; 28(10): 2237-2251.
 42. Laborda E, Mazagova M, Shao S, Wang X, Quirino H, Woods AK, et al. "Development of a chimeric antigen receptor targeting C-type lectin-like molecule-1 for human acute myeloid leukemia." *International journal of molecular sciences*, 2017; 18(11): 2259.
 43. Zhang H, Gan WT, Hao WG, Wang PF, Li ZY, Chang LJ. "Successful anti-CLL1 CAR T-cell therapy in secondary acute myeloid leukemia." *Frontiers in Oncology*, 2020; 10: 685.
 44. Loff S, Dietrich J, Meyer JE, Riewaldt J, Spehr J, von Bonin M, et al. "Rapidly switchable universal CAR-T cells for treatment of CD123-positive leukemia." *Molecular Therapy-Oncolytics*, 2020; 17: 408-420.
 45. El Khawanky N, Hughes A, Yu W, Myburgh R, Matschulla T, Taromi S, et al. "Demethylating therapy increases anti-CD123 CAR T cell cytotoxicity against acute myeloid leukemia." *Nature communications*. 2021; 12(1): 1-20.
 46. Gomes-Silva D, Atilla E, Atilla PA, Mo F, Tashiro H, Srinivasan M, et al. "CD7 CAR T cells for the therapy of acute myeloid leukemia." *Molecular Therapy*, 2019; 27(1): 272-280.
 47. Gill, Saar I. "How close are we to CAR T-cell therapy for AML?" *Best practice & research Clinical haematology*, 2019; 32(4): 101104.
 48. Müller LM, Holmes M, Michael JL, Scott GB, West EJ, Scott KJ, et al. "Plasmacytoid dendritic cells orchestrate innate and adaptive anti-tumor immunity induced by oncolytic coxsackievirus A21." *Journal for immunotherapy of cancer*, 2019; 7(1): 1-6.
 49. Bais Swarna, et al. "Oncolytic virotherapy for hematological malignancies." *Advances in virology* 2012, 2012.
 50. Wu Ming, et al. "Therapeutic cancer vaccine therapy for acute myeloid leukemia." *Immunotherapy* 2021; 13(10): 863-877.
 51. Kerage D, Soon MS, Doff BL, Kobayashi T, Nissen MD, Lam PY, et al. "Therapeutic vaccination with 4-1BB co-stimulation eradicates mouse acute myeloid leukemia." *OncoImmunology*, 2018; 7(10): e1486952.
 52. Liu Yuxin, et al. "Immunotherapy in acute myeloid leukemia and myelodysplastic syndromes: The dawn of a new era?" *Blood reviews*, 2019; 34: 67-83.
 53. Minev BR, Lander E, Feller JF, Berman M, Greenwood BM, Minev I, et al. "First-in-human study of TK-positive oncolytic vaccinia virus delivered by adipose stromal vascular fraction cells." *Journal of translational medicine*, 2019; 17(1): 1-5.
 54. Anguille Sébastien, et al. "Interleukin-15 dendritic cells as vaccine candidates for cancer immunotherapy." *Human vaccines & immunotherapeutics*; 2013; 9(9): 1956-1961.
 55. Maurer S, Salih HR, Smirnow I, Lauer UM, Berchtold S. "Suicide gene armed measles vaccine virus for the treatment of AML." *International journal of oncology*, 2019; 55(2): 347-358.
 56. Van Acker HH, Anguille S, De Reu H, Berneman ZN, Smits EL, Van Tendeloo VF. "Interleukin-15-cultured dendritic cells enhance anti-tumor gamma delta T cell functions through IL-15 secretion." *Frontiers in immunology*, 2018; 9: 658.
 57. Sendker Stephanie, Dirk Reinhardt, Naghmeh Niktoresh. "Redirecting the immune microenvironment in acute myeloid leukemia." *Cancers*, 2021; 13(6): 1423.
 58. Shi Y, Dincheva-Vogel L, Ayemoba CE, Fung JP, Bergamaschi C, Pavlakis GN, et al. "IL-15/IL-15R α /CD80-expressing AML cell vaccines eradicate minimal residual disease in leukemic mice." *Blood advances*, 2018; 2(22): 3177-3192.
 59. Subklewe Marion, et al. "New generation dendritic cell vaccine for immunotherapy of acute myeloid leukemia." *Cancer Immunology, Immunotherapy*, 2014; 63(10): 1093-1103.
 60. Anguille S, Van de Velde AL, Smits EL, Van Tendeloo VF, Juliusson G, Cools N, et al. "Dendritic cell vaccination as postremission treatment to prevent or delay relapse in acute myeloid leukemia." *Blood, The Journal of the American Society of Hematology*, 2017; 130(15): 1713-1721.
 61. Maslak PG, Dao T, Bernal Y, Chanel SM, Zhang R, Frattini M, et al. "Phase 2 trial of a multivalent WT1 peptide vaccine (galinpepimut-S) in acute myeloid leukemia." *Blood advances*, 2018; 2(3): 224-234.
 62. Van Acker, Heleen H, et al. "Dendritic cell-based immunotherapy of acute myeloid leukemia." *Journal of clinical medicine*, 2019; 8(5): 579.
 63. van de Loosdrecht AA, van Wetering S, Santegoets SJ, Singh SK, Eeltingk CM, den Hartog Y, et al. "A novel allogeneic off-the-shelf dendritic cell vaccine for post-remission treatment of elderly patients with acute myeloid leukemia." *Cancer Immunology*,

- Immunotherapy, 2018; 67(10): 1505-1518.
64. Wang X, Yang Y, Wang N, Wu X, Xu J, Zhou Y, et al. "Mesenchymal stem cell carriers enhance antitumor efficacy induced by oncolytic reovirus in acute myeloid leukemia." *International Immunopharmacology*, 2021; 94: 107437.
 65. Hadrys Agata, et al. "Mesenchymal stem cells as carriers for systemic delivery of oncolytic viruses." *European journal of pharmacology*, 2020; 874: 172991.
 66. Johnson DT, Zhou J, Kroll AV, Fang RH, Yan M, Xiao C, et al. Acute myeloid leukemia cell membrane-coated nanoparticles for cancer vaccination immunotherapy." *Leukemia*, 2021: 1-2.
 67. Shetab Boushehri, Maryam A, Alf Lamprecht. "TLR4-based immunotherapeutics in cancer: a review of the achievements and shortcomings." *Molecular pharmaceutics*, 2018; 15(11): 4777-4800.
 68. Zhong G, Jin G, Zeng W, Yu C, Li Y, Zhou J, et al. "Conjugation of TLR7 Agonist Combined with Demethylation Treatment Improves Whole-Cell Tumor Vaccine Potency in Acute Myeloid Leukemia." *International Journal of Medical Sciences*, 2020; 17(15): 2346.
 69. Melzer MK, Zeitlinger L, Mall S, Steiger K, Schmid RM, Ebert O, et al. "Enhanced safety and efficacy of oncolytic VSV therapy by combination with T cell receptor transgenic T cells as carriers." *Molecular Therapy-Oncolytics*. 2019; 12: 26-40.
 70. Mi Yu, et al. "Emerging nano-/microapproaches for cancer immunotherapy." *Advanced Science*, 2019; 6(6): 1801847.
 71. Zhang C, Wang XQ, Zhang RL, Liu F, Wang Y, Yan ZL, et al. Donor-derived CD19 CAR-T cell therapy of relapse of CD19-positive B-ALL post allotransplant. *Leukemia*, 2021; 35(6): 1563-1570.
 72. Sheth Vipul S, Jordan Gauthier. "Taming the beast: CRS and ICANS after CAR T-cell therapy for ALL." *Bone marrow transplantation*, 2021; 56(3): 552-566.
 73. Cappell KM, Sherry RM, Yang JC, Goff SL, Vanasse DA, McIntyre L, et al. "Long-term follow-up of anti-CD19 chimeric antigen receptor T-cell therapy." *Journal of Clinical Oncology*, 2020; 38(32): 3805-3815.
 74. Santomaso BD, Park JH, Salloum D, Riviere I, Flynn J, Mead E, et al. "Clinical and biological correlates of neurotoxicity associated with CAR T-cell therapy in patients with B-cell acute lymphoblastic leukemia." *Cancer discovery*, 2018; 8(8): 958-971.
 75. Korell F, Schubert ML, Sauer T, Schmitt A, Derigs P, Weber TF, et al. "Infection Complications after Lymphodepletion and Dosing of Chimeric Antigen Receptor T (CAR-T) Cell Therapy in Patients with Relapsed/Refractory Acute Lymphoblastic Leukemia or B Cell Non-Hodgkin Lymphoma." *Cancers*, 2021; 13(7): 1684.
 76. Zhang C, Wang XQ, Zhang RL, Liu F, Wang Y, Yan ZL, et al. "Donor-derived CD19 CAR-T cell therapy of relapse of CD19-positive B-ALL post allotransplant." *Leukemia*, 2021; 35(6): 1563-1570.
 77. Pillai V, Muralidharan K, Meng W, Bagashev A, Oldridge DA, Rosenthal J, et al. "CAR T-cell therapy is effective for CD19-dim B-lymphoblastic leukemia but is impacted by prior blinatumomab therapy." *Blood advances*, 2019; 3(22): 3539-3549.
 78. Zhao Juanjuan, Yongping Song, Delong Liu. "Recent advances on blinatumomab for acute lymphoblastic leukemia." *Experimental Hematology & Oncology*, 2019; 8(1): 1-8.
 79. Amrolia PJ, Wynn R, Hough RE, Vora A, Bonney D, Veys P, et al. "Phase I study of AUTO3, a bicistronic chimeric antigen receptor (CAR) T-cell therapy targeting CD19 and CD22, in pediatric patients with relapsed/refractory B-cell acute lymphoblastic leukemia (r/r B-ALL): Amelia Study." *Blood*, 2019; 134: 2620.
 80. Liu S, Deng B, Yin Z, Lin Y, An L, Liu D, et al. "Combination of CD19 and CD22 CAR-T cell therapy in relapsed B-cell acute lymphoblastic leukemia after allogeneic transplantation" *American Journal of Hematology*, 2021; 96(6): 671-679.
 81. Shah BD, Bishop MR, Oluwole OO, Logan AC, Baer MR, Donnellan WB, et al. "KTE-X19 anti-CD19 CAR T-cell therapy in adult relapsed/refractory acute lymphoblastic leukemia: ZUMA-3 phase 1 results." *Blood, The Journal of the American Society of Hematology*, 2021; 138(1): 11-22.
 82. Li S, Zhang J, Wang M, Fu G, Li Y, Pei LI, et al. "Treatment of acute lymphoblastic leukaemia with the second generation of CD19 CAR-T containing either CD 28 or 4-1 BB." *British journal of haematology*, 2018; 181(3): 360-371.
 83. Li Z, Chen X, Liu L, Zhou M, Zhou G, Liu T. "Development of the T-ALLiPSC-based Therapeutic Cancer Vaccines for T-cell acute Lymphoblastic Leukemia."
 84. Rossig C, Pule M, Altwater B, Saiagh S, Wright G, Ghorashian S, et al. "Vaccination to improve the persistence of CD19CAR gene-modified T cells in relapsed pediatric acute lymphoblastic leukemia." *Leukemia*, 2017; 31(5): 1087-1095.
 85. Wu M, Zhang L, Zhang H, Ning J, Tu S, He Y, et al. "CD19 chimeric antigen receptor–redirected T cells combined with epidermal growth factor receptor pathway substrate 8 peptide–derived dendritic cell vaccine in leukemia." *Cytotherapy*. 2019; 21(6): 659-670.
 86. Rengifo, Andrés Felipe Chamorro, et al. "A new and efficient carboxymethyl-hexanoyl chitosan/dodecyl sulfate nanocarrier for a pyrazoline with antileukemic activity." *Materials Science and Engineering: C*, 2019; 105: 110051.
 87. Kumar, Lalit, et al. "Nanofiber as a novel vehicle for transdermal delivery of therapeutic agents: challenges and opportunities." *Future Journal of Pharmaceutical Sciences*, 2021; 7(1): 1-17.
 88. Wu, Xiaoyan, et al. "Cooperation of IRAK1/4 inhibitor and ABT-737 in nanoparticles for synergistic therapy of T cell acute lymphoblastic leukemia." *International Journal of Nanomedicine*, 2017; 12: 8025.
 89. Bayat Narges, et al. "Thiol-reactive star polymers functionalized with short ethoxy-containing moieties exhibit enhanced uptake in acute lymphoblastic leukemia cells." *International Journal of Nanomedicine*, 2019; 14: 9795.
 90. Tinoco Ana, et al. "BSA/ASN/Pol407 nanoparticles for acute lymphoblastic leukemia treatment." *Biochemical Engineering Journal*, 2019; 141: 80-88.
 91. Nunes João CF, et al. "Recent strategies and applications for L-asparaginase confinement." *Molecules*, 2020; 25(24): 5827.
 92. Gan Donghui, et al. "Doxorubicin/Nucleophosmin Binding Protein-Conjugated Nanoparticle Enhances Anti-leukemia Activity in Acute Lymphoblastic Leukemia Cells in vitro and in vivo." *Frontiers in pharmacology*, 2021; 12: 1255.
 93. Li Shanshan, et al. "Anti-CD19 mAb modified mesoporous titanium dioxide as exclusively targeting vector for efficient B-lymphoblastic leukemia therapy." *Journal of Pharmaceutical Sciences*, 2021; 110(7): 2733-2742.
 94. Bailey Katharine E, et al. "The Relevance of CD20 Targeting of Oncolytic Measles Virus in the Treatment of Adult Acute Lymphoblastic Leukemia." *Blood*, 2017; 130: 3891.
 95. Castleton Anna, et al. "Human mesenchymal stromal cells deliver systemic oncolytic measles virus to treat acute lymphoblastic leukemia in the presence of humoral immunity." *Blood, The Journal of the American Society of Hematology*, 2014; 123(9): 1327-1335.
 96. Chua LL, Rajasuriar R, Lim YA, Woo YL, Ariffin H. "Temporal changes in gut microbiota profile in children with acute lymphoblastic leukemia prior to commencement-, during-, and post-cessation of chemotherapy." *BMC cancer*, 2020; 20(1): 1-1.
 97. De Pietri S, Ingham AC, Frandsen TL, Rathe M, Krych L, Castro-Mejia JL, et al. "Gastrointestinal toxicity during induction treatment for childhood acute lymphoblastic leukemia: the impact of the gut microbiota." *International journal of cancer*, 2020; 147(7): 1953-1962.
 98. Rattanathammethee T, Tuitemwong P, Thiennimitr P, Sarichai P, Na Pombejra S, Piriyaakuntorn P, et al. "Gut microbiota profiles of treatment-naïve adult acute myeloid leukemia patients with neutropenic fever during intensive chemotherapy." *PloS one*, 2020; 15(10): e0236460.
 99. Hueso T, Ekpe K, Mayeur C, Gatse A, Joncquel-Chevallier Curt M, Gricourt G, et al. "Impact and consequences of intensive chemotherapy on intestinal barrier and microbiota in acute myeloid leukemia: the role of mucosal strengthening." *Gut microbes*, 2020; 12(1): 1800897
 100. Song Y, Gyarmati P. "Microbiota changes in a pediatric acute lymphocytic leukemia mouse model." *MicrobiologyOpen*, 2020; 9(3): e982.
 101. Rashidi A, Ebadi M, Rehman TU, Elhusseini H, Nalluri H, Kaiser T, et al. Altered "microbiota-host metabolic cross talk preceding neutropenic fever in patients with acute leukemia." *Blood*

- Advances. 2021; 5(20): 3937-3950.
102. Malard F, Vekhoff A, Lapusan S, Isnard F, D'incan-Corda E, Rey J, et al. "Gut microbiota diversity after autologous fecal microbiota transfer in acute myeloid leukemia patients." *Nature communications*. 2021; 12(1): 1-1.
103. Lee MS, Ramakrishna B, Moss AC, Gold HS, Branch-Elliman W. "Successful treatment of fulminant *Clostridioides difficile* infection with emergent fecal microbiota transplantation in a patient with acute myeloid leukemia and prolonged, severe neutropenia." *Transplant Infectious Disease*, 2020; 22(1): e13216.
104. Shen Z, Gu X, Cao H, Mao W, Yang L, He M, et al. "Characterization of microbiota in acute leukemia patients following successful remission induction chemotherapy without antimicrobial prophylaxis." *International Microbiology*, 2021; 24(2): 263-273.